

# Formation of multi-solitons and vortex bright solitons in Bose-condensed alkali-metal atoms

Luca Salasnich

*INFN-CNR, Unità di Milano Università,  
Dipartimento di Fisica, Università di Milano,  
Via Celoria 16, Milano 20133, Italy  
salasnich@mi.infn.it*

## Abstract

The formation of multi-soliton configurations in Bose-condensed alkali-metal atoms is analyzed by using the nonpolynomial Schrödinger equation. A train of bright solitons is obtained from an axially homogeneous Bose-Einstein condensate by a sudden change of the scattering length from repulsive to attractive. We derive an analytical expression for the number of bright solitons generated by using this mechanism. The formula generalizes a previous formula obtained with the 1D Gross-Pitaevskii equation. In the second part we consider vortex bright solitons, namely cigar-shaped bright solitons with a nonzero angular quantum number  $k$  along the axial direction. By using a variational approach we determine the shape of vortex bright solitons, showing that the critical number of atoms for the collapse of the vortex soliton increases with a larger  $k$ . Finally we calculate monopole and quadrupole collective oscillations of these vortex bright solitons.

## I. INTRODUCTION

In the last years we have investigated various dynamical properties of pure Bose-Einstein condensates (BECs) made of alkali-metal atoms at ultra-low temperatures [1]. The recent experimental observation of dark [2] and bright [3,4] solitons in Bose-Einstein condensates (BECs) has focused our interest on BEC bright solitons. In particular, by using the non-polynomial Schrödinger equation (NPSE), an effective equation derived from the 3D Gross-Pitaevskii equation (GPE) [5], we have studied the stability properties of bright solitons, determining the critical strength beyond which bright solitons collapse [6]. This collapse is a direct consequence of the non-constant size of the soliton in the transverse direction, as found by solving the 3D GPE and NPSE. Instead, the 1D GPE, for which the transverse size of the soliton is constant, admits only stable bright solitons [6].

In this paper we analyze the formation of bright solitons in a Bose-Einstein condensate of  $^7\text{Li}$  atoms induced by a sudden change in the sign of the scattering length from positive to negative, as reported in a recent experiment [4]. In a previous paper [7] we have shown that a number of bright solitons is produced and this can be interpreted in terms of the modulational instability of the time-dependent macroscopic wave function of the Bose condensate. From the 1D GPE we have derived a simple formula for the number of solitons that is in good agreement with the numerical results. Here we generalize such a formula taking into account the non-constant size of the BEC in the transverse direction. Then we investigate the properties of vortex bright solitons, which are bright solitons with a finite quantum number of circulation around the longitudinal axis.

## II. FORMATION OF BEC BRIGHT SOLITONS

At zero temperature the macroscopic wave function of a BEC made of  $N$  dilute  $^7\text{Li}$  atoms can be modeled by the 3D GPE. For a dilute gas the inter-atomic strength depends on the single parameter: the s-wave scattering length  $a_s$ . Note that only for  $a_s < 0$  there

exist bright solitons, which are shape-invariant waves traveling without spreading due to the interplay between the repulsive kinetic energy and the attractive inter-atomic energy.

Following the experiment of Strecker *et al.* [4], the trapping potential of the BEC is modeled by an anisotropic harmonic potential with transverse frequency  $\omega_{\perp} = 2\pi \times 800$  Hz and longitudinal frequency  $\omega_z = 2\pi \times 4$  Hz. In addition there is a box optical potential that initially confines the condensate in the longitudinal direction.

By using 3D GPE we first calculate the ground-state wave function of the condensate with a positive scattering length  $a_s = 100a_B$ , where  $a_B = 0.53\text{\AA}$  is the Bohr radius. We choose a condensate with longitudinal width  $L = 284.4 \mu\text{m}$  and  $N = 10^4$  atoms. Then the ground-state wave function is inserted as initial condition for the time-evolution of the 3D GPE with  $a_s = -3a_B$ . We observe that a three-body dissipative term has been added to the 3D GPE. This term is important mainly during the collapse because it fixes the critical density at which the compression ceases (see also [7]).

In Figure 1 we plot the probability density in the longitudinal direction  $\rho(z)$  of the evolving wave-function. The Bose-Einstein condensate shows the formation of 5 peaks. It is remarkable to stress that at the first instants the two external peaks become very high but the collapse is avoided by the dissipative term that quickly reduces their density. We have verified that without the dissipative term the Bose condensate collapses. The dissipative term does not alter the number of peaks but only their density. After the formation, the peaks start to separate each other showing a repulsive force between them. Eventually each peak evolves with small shape oscillations but without appreciable dispersion.

The formation of these solitonic peaks can be explained as due to MI of the time-dependent wave function of the BEC, driven by imaginary Bogoliubov excitations [7]. Neglecting the dissipative term, in the 1D limit of very long cigar-shaped condensate, the Bogoliubov longitudinal elementary excitations  $\epsilon_k$  of the axially homogeneous BEC can be obtained from the 1D GPE. They are given by

$$\epsilon_k = \sqrt{\frac{\hbar^2 k^2}{2m} \left( \frac{\hbar^2 k^2}{2m} + 2g_{1d}n \right)}, \quad (1)$$

where  $n = N/L$  is the linear axial density and  $g_{1d} = 2(\hbar\omega_{\perp})a_s$  is the 1D interaction strength. By suddenly changing the scattering length  $a_s$  to a negative value, the excitations frequencies corresponding to  $k < k_c = (16\pi|g_{1d}|n)^{1/2}$  become imaginary and, as a result, small perturbations grow exponentially in time. It is easy to find that the maximum rate of growth is at  $k_0 = k_c/\sqrt{2}$ . The wavelength of this mode is  $\lambda_0 = 2\pi/k_0$  and the ratio  $L/\lambda_0$  gives an estimate of the number  $N_s$  of bright solitons which are generated:

$$N_s = \frac{\sqrt{N|a_s|L}}{\pi a_{\perp}}, \quad (2)$$

with  $a_{\perp} = (\hbar/(m\omega_{\perp}))^{1/2}$ . The predicted number  $N_s$  of solitons is in very good agreement with the numerical results of Figure 1 (see also [7]).

The Eq. (2) is valid in the 1D limit but can be extended to a generic 3D case by using NPSE instead of 1D GPE. In general, the Bogoliubov excitations can be written as

$$\epsilon_k = \sqrt{\frac{\hbar^2 k^2}{2m} \left( \frac{\hbar^2 k^2}{2m} + 2c^2 \right)}, \quad (3)$$

where  $c$  is the first sound velocity in the longitudinal direction. The sound velocity  $c$  satisfies the equation

$$c = \sqrt{n \frac{\partial \mu}{\partial n}}, \quad (4)$$

where  $\mu$  is the chemical potential of the Bose condensate. The chemical potential derived from 1D GPE is  $\mu = g_{1d}n + \hbar\omega_{\perp}$ , while using NPSE one finds

$$\mu = \frac{g_{1d}n}{\sqrt{1+2a_s n}} + \frac{\hbar\omega_{\perp}}{2} \left( \frac{1}{\sqrt{1+2a_s n}} + \sqrt{1+2a_s n} \right). \quad (5)$$

This chemical potential reduces to the 1D GPE one in the 1D limit, i.e. for  $2a_s n \ll 1$ . Moreover, if  $a_s < 0$  the chemical potential  $\mu$  diverges at  $2|a_s|n = 1$ , as a consequence of the collapse on the condensate [8]. In conclusion, NPSE tells us that the number  $N_s$  of bright solitons obtained via modulational instability is given by

$$N_s = \frac{g\sqrt{2}}{4\pi} \frac{\sqrt{4-3g\rho}}{\sqrt{g\rho}(1-g\rho)^{3/4}}, \quad (6)$$

where  $g = 2|a_s|/a_\perp$  is the adimensional inter-atomic strength and  $\rho = na_\perp$  is the adimensional axial density of the condensate. In Figure 2 we plot  $N_s$  as a function of  $g\rho$  for a fixed value of  $\rho$ . For small values of  $g\rho$  1D GPE and NPSE give the same results but approaching  $g\rho = 1$  the results are strongly different. In fact, NPSE gives a divergent number of solitons at  $g\rho = 1$ . As previously stated, only NPSE takes into account the non-constant width of the BEC in the transverse direction. Such a width shrinks to zero for a large attractive inter-atomic interaction.

### III. VORTEX BRIGHT SOLITONS

The 3D GPE of a Bose-Einstein condensate of  $N$  atoms confined in the transverse direction by a harmonic potential of frequency  $\omega_\perp$  and with an attractive inter-atomic interaction ( $a_s < 0$ ) can be derived by the following scaled GPE Lagrangian

$$L = \psi^* \left[ i \frac{\partial}{\partial t} + \frac{1}{2} \nabla^2 - \frac{1}{2} (x^2 + y^2) - 2\pi\gamma |\psi|^2 \right] \psi \quad (7)$$

where  $\psi$  is the macroscopic wave function of the condensate and  $\gamma = Na_s/a_\perp$ , with  $a_\perp = (\hbar/(m\omega_\perp))^{1/2}$ . Note that in the GPE Lagrangian lengths are in units  $a_\perp$ , time in units  $\omega_\perp^{-1}$ , action in units  $\hbar$  and energy in units  $\hbar\omega_\perp$ .

Using the cylindrical coordinates  $r = (x^2 + y^2)^{1/2}$ ,  $\theta = \arctg(y/x)$  and  $z$ , we write the trial wave function of the vortex bright soliton as

$$\psi(r, z, \theta, t) = r^k e^{ik\theta} \exp \left\{ -\frac{r^2}{2\sigma(t)^2} - \frac{z^2}{2\eta(t)^2} + i\alpha(t)r^2 + i\beta(t)z^2 \right\}, \quad (8)$$

where  $k$  is the vortex quantum number and  $\sigma$ ,  $\eta$ ,  $\alpha$  and  $\beta$  are the time-dependent variational parameters. In [9] we have used this trial wave function to study vortex states with positive scattering length in a harmonic potential. Here we consider a negative scattering length. Moreover, in this case the (harmonic) confinement is only in the transverse direction. By inserting the trial wave function in the GPE Lagrangian, after spatial integration, one gets an effective Lagrangian depending on the variational parameters. By using the Euler-Lagrange

equations one realizes that the time dependence of the phase parameters  $\alpha$  and  $\beta$  is fully determined by that of  $\sigma$  and  $\eta$  (see also [9]). Thus, one obtains a new effective Lagrangian depending on  $\sigma$  and  $\eta$  only. It is given by

$$L_{eff} = (k+1)\dot{\sigma}^2 + \frac{1}{2}\dot{\eta}^2 - (k+1)\sigma^2 - \frac{(k+1)}{\sigma^2} - \frac{1}{2\eta^2} - \sqrt{\frac{2}{\pi}} \frac{(2k)!}{2^{2k}(k!)^2} \frac{\gamma}{\sigma^2\eta}. \quad (9)$$

In the case  $k = 0$  this effective Lagrangian reduces to the one we have successfully used to analyze the collective oscillations of the bright soliton in [6]. In fact, the stationary solutions of the Euler-Lagrange equations of  $L_{eff}$  give the widths of the bright soliton ( $k = 0$ ) and the widths of the vortex bright soliton ( $k \neq 0$ ).

In Figure 3 we plot the transverse radial width  $\sigma$  and the longitudinal axial width  $\eta$  of the vortex bright soliton as a function of the inter-atomic strength  $\gamma$ . For values of  $\gamma$  close to zero the axial width  $\eta$  is very large with respect to the radial width  $\sigma$ : the soliton is a strongly elongated cigar. By decreasing  $\gamma$  the soliton anisotropy is reduced up to the collapse, i.e. the critical point at which the width parameters are no more minima of the GPE potential energy. It is interesting to observe that the critical  $\gamma$  changes with the vortex quantum number  $k$ :  $\gamma_c = -0.78$  for  $k = 0$ ,  $\gamma_c = -2.20$  for  $k = 1$ , and  $\gamma_c = -3.59$  for  $k = 2$ . This means that it is possible to avoid the collapse by creating vortex bright solitons with a large vortex quantum number  $k$  (see also the recent numerical results [10] of Adhikari). Another remarkable effect shown in Figure 3 for  $k \neq 0$  is the crossing of the widths by decreasing  $\gamma$ : the vortex soliton becomes slightly disc-shaped.

In the lower part of Figure 3 we plot the frequencies  $\Omega_1$  and  $\Omega_2$  of the collective oscillations of the vortex bright soliton along the axial and transverse directions. In cigar-shaped condensates  $\Omega_1$  is called monopole frequency and  $\Omega_2$  is called quadrupole frequency [9]. These frequencies are obtained from the effective Lagrangian  $L_{eff}$  linearizing around the minima of the potential energy (equilibrium solution). The transverse frequency  $\Omega_2$  remains practically constant while the axial frequency  $\Omega_1$  grows by decreasing  $\gamma$  but goes to zero at the critical point, beyond which the frequencies are complex and the equilibrium solution becomes unstable. Note that for a fixed  $\gamma$ , the value of  $\Omega_1$  strongly depends on  $k$ .

## IV. CONCLUSIONS

In this paper we have shown that a train of bright solitons can be created in a Bose-Einstein condensate by a sudden change of the scattering length from positive to negative. The number of bright solitons can be predicted by using an analytical formula we have derived from the nonpolynomial Schrödinger equation. We have also investigated vortex bright solitons, which are bright solitons with a finite angular momentum along the longitudinal axis. Our results suggest that vortex bright solitons can be created with a larger number of atoms than bright solitons without vorticity. Moreover, our calculations suggest that vortex bright solitons can be experimentally detected by analyzing their collective oscillations.

## REFERENCES

- [1] L. Salasnich, Int. J. Mod. Phys. B **15**, 1253 (2001); L. Salasnich, Int. J. Mod. Phys. B **16**, 2185 (2002); L. Salasnich, Nuovo Cimento B **114**, 637 (2002); L. Salasnich, Laser Phys. **13**, 547 (2003);
- [2] S. Burger et al., Phys. Rev. Lett. **83**, 5198 (1999); J. Denschlag *et al.*, Science **287**, 97 (2000).
- [3] L. Khaykovich *et al.*, Science **296**, 1290 (2002).
- [4] K.E. Strecker *et al.*, Nature **417**, 150 (2002).
- [5] L. Salasnich, Laser Phys. **12**, 198 (2002); L. Salasnich, A. Parola and L. Reatto, Phys. Rev. A **65**, 043614 (2002); L. Salasnich, A. Parola and L. Reatto, J. Phys. B (Atom. Mol.) **35**, 3205 (2002). L. Salasnich, Laser Phys. **13**, 543 (2003).
- [6] L. Salasnich, A. Parola and L. Reatto, Phys. Rev. A **66**, 043603 (2002); L. Salasnich, Progr. Theor. Phys. Suppl. **150** (2003).
- [7] L. Salasnich, A. Parola and L. Reatto, cond-mat/0307206, to be published in Phys. Rev. Lett. (2003).
- [8] L. Salasnich, A. Parola and L. Reatto, in preparation.
- [9] L. Salasnich, Int. J. Mod. Phys. B **14**, 1 (2000).
- [10] S.K. Adhikari, Phys. Rev. E **65**, 016703 (2002).



## FIGURE CAPTIONS

**Figure 1:** Axial density profile  $\rho(z)$  of the Bose-Einstein condensate made of  $10^4$   $^7\text{Li}$  atoms obtained by solving 3D GPE. For  $t < 0$  the scattering length is  $a_s = 100a_B$ , while for  $t \geq 0$  we set  $a_s = -3a_B$  with  $a_B$  the Bohr radius. Length  $z$  in units  $a_z = (\hbar/m\omega_z)^{1/2}$ , density  $\rho$  in units  $1/a_z$  and time  $t$  in units  $\omega_z^{-1}$ . Adapted from [7].

**Figure 2:** Number  $N_s$  of bright solitons created via modulational instability.  $\rho = Na_\perp/L$  is the initial (adimensional) axial density of the axially homogeneous Bose condensate.  $g = 2|a_s|/a_\perp$  is the inter-atomic strength with  $a_\perp = (\hbar/m\omega_\perp)^{1/2}$  and  $a_s < 0$  the final scattering length.

**Figure 3:** Top: transverse width  $\sigma$  and axial width  $\eta$  of the vortex bright soliton as a function of the interaction strength  $\gamma = Na_s/a_\perp$ . Bottom: monopole frequency  $\Omega_1$  and quadrupole frequency  $\Omega_2$  of the vortex bright soliton as a function of the interaction strength  $\gamma$ . Lengths in units  $a_\perp$  and frequencies in units  $\omega_\perp$ .  $k$  is the vortex quantum number.